

Misère Quotients for Impartial Games

SUPPLEMENTARY MATERIAL

Thane E. Plambeck
2341 Tasso St.
Palo Alto, CA 94301

Aaron N. Siegel
Institute for Advanced Study
1 Einstein Drive
Princeton, NJ 08540

February 1, 2008

These are the supplementary appendices to the paper, Misère quotients for impartial games [5]. If you have not read the actual paper, you should do so *before* reading this supplement, or else it will not make much sense!

Appendix B gives detailed solutions to many of the octal games discussed in the paper, and Appendix C describes the algorithms used to compute most of our solutions.

B Solutions in Detail

This appendix contains detailed solutions to many of the games discussed in Appendix A.

Figure 1 summarizes the status of every octal game with at most three code digits. For each game Γ , the chart indicates whether Γ is tame or wild, and whether its normal- and/or misère-play solution is known.

Figures 2 and 4 present complete solutions to wild two- and three-digit octal games with relatively simple misère quotients and pretending functions. Figure 3 does the same for wild four-digit quaternaries. Finally, Figures 5, 6 and 7 present the solutions to **0.115**, **0.152** and **0.77**, respectively. Note that $\mathcal{Q}(\mathbf{0.15}) \cong \mathcal{Q}(\mathbf{0.115})$. The solution to **0.644** is omitted due to its size; see [4].

$d_1 d_2$	d_3								$d_1 d_2$	d_3							
	0	1	2	3	4	5	6	7		0	1	2	3	4	5	6	7
.00	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	—	—	.44	$\ominus$	$\ominus$	$\ominus$	$\ominus$	—	—	—	—
.01	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	—	$\star$	.45	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$
.02	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	—	—	.46	$\ominus$	$\ominus$	$\ominus$	$\ominus$	—	—	—	—
.03	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	—	—	.47	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$
.04	—	—	—	$\star$	$\ominus$	$\star$	$\ominus$	$\star$	.50	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$
.05	$\oplus$	$\star$	$\oplus$	$\star$	$\star$	$\star$	—	$\star$	.51	$\oplus$	$\oplus$	$\ominus$	$\oplus$	$\oplus$	$\star$	$\star$	$\star$
.06	—	—	—	—	—	—	—	—	.52	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\star$	$\star$	$\star$	$\star$
.07	$\star$	$\star$	$\star$	$\star$	$\ominus$	$\ominus$	$\ominus$	$\ominus$	.53	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\star$	$\star$	$\star$	$\star$
.10	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\star$	$\oplus$	$\star$	$\oplus$	.54	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$
.11	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	$\ominus$	$\star$	$\star$	.55	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\star$	$\star$	$\star$
.12	$\oplus$	$\oplus$	$\oplus$	$\ominus$	$\star$	—	—	$\star$	.56	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	—	—
.13	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\star$	—	—	$\star$	.57	$\ominus$	$\ominus$	$\oplus$	$\oplus$	$\star$	$\star$	$\star$	$\star$
.14	—	$\star$	—	—	$\ominus$	$\star$	—	$\oplus$	.60	—	—	$\oplus$	$\oplus$	—	—	—	—
.15	$\ominus$	$\oplus$	$\ominus$	$\ominus$	$\star$	$\oplus$	$\oplus$	$\ominus$	.61	—	—	$\oplus$	$\oplus$	—	—	—	—
.16	—	—	—	—	—	$\star$	—	—	.62	—	—	$\oplus$	$\oplus$	—	—	—	—
.17	$\star$	$\star$	—	$\star$	—	$\star$	$\star$	$\star$	.63	—	—	$\oplus$	$\oplus$	—	—	—	—
.20	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	—	—	.64	—	—	—	—	$\ominus$	$\ominus$	$\ominus$	$\ominus$
.21	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	—	—	.65	—	—	—	—	$\ominus$	$\ominus$	$\ominus$	$\ominus$
.22	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	$\star$	$\star$	.66	—	—	—	—	$\ominus$	$\ominus$	$\ominus$	$\ominus$
.23	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	$\star$	$\star$	.67	—	—	—	—	$\ominus$	$\ominus$	$\ominus$	$\ominus$
.24	$\oplus$	$\ominus$	$\oplus$	$\ominus$	—	—	—	—	.70	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$
.25	$\oplus$	$\ominus$	$\oplus$	$\ominus$	—	—	—	—	.71	$\ominus$	$\ominus$	$\ominus$	$\ominus$	$\star$	$\star$	$\ominus$	$\ominus$
.26	$\star$	$\star$	$\star$	$\star$	—	—	—	—	.72	$\ominus$	$\ominus$	$\ominus$	$\ominus$	$\ominus$	$\ominus$	$\ominus$	$\ominus$
.27	$\star$	$\star$	$\star$	$\star$	—	—	—	—	.73	$\oplus$	$\oplus$	$\star$	$\star$	$\star$	$\star$	$\star$	$\star$
.30	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	.74	—	—	—	—	—	—	—	—
.31	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	$\ominus$	$\star$	$\ominus$	.75	$\ominus$	$\ominus$	$\ominus$	$\ominus$	$\star$	$\star$	$\star$	$\star$
.32	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	—	$\star$	$\star$	.76	—	—	—	—	—	—	—	—
.33	$\oplus$	$\oplus$	$\oplus$	$\oplus$	—	$\star$	$\star$	$\star$	.77	$\ominus$	$\ominus$	$\oplus$	$\oplus$	—	—	—	—
.34	$\ominus$	$\ominus$	—	—	—	—	—	—	4.0	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\star$	$\star$	$\star$
.35	$\star$	$\ominus$	$\oplus$	$\oplus$	—	$\star$	$\star$	$\star$	4.1	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\star$	$\star$	$\star$
.36	—	—	—	—	—	—	—	—	4.2	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\star$	$\star$	$\star$
.37	—	—	$\oplus$	$\oplus$	—	$\star$	$\star$	—	4.3	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$	$\oplus$
.40	$\star$	$\star$	$\star$	$\star$	—	—	—	—	4.4	$\ominus$	$\ominus$	$\ominus$	$\ominus$	—	—	—	—
.41	$\star$	$\star$	$\star$	$\star$	—	—	—	—	4.5	$\oplus$	$\oplus$	$\star$	$\star$	$\oplus$	$\oplus$	$\ominus$	$\ominus$
.42	$\star$	$\star$	$\star$	$\star$	—	—	—	—	4.6	$\ominus$	$\ominus$	$\ominus$	$\ominus$	—	—	—	—
.43	$\star$	$\star$	$\star$	$\star$	—	—	—	—	4.7	$\ominus$	$\ominus$	$\oplus$	$\oplus$	$\ominus$	$\ominus$	$\star$	$\star$

- $\oplus$ Tame; solution is known in both normal and misère play.
- $\ominus$ Wild; solution is known in both normal and misère play. See [5] for details.
- Wild; solution is unknown in either normal or misère play. For details on two-digit octals, see [5]; for three-digit octals, see [4].
- $\star$ Wild; solution is known in normal play, but not in misère play. These are the most promising candidates for further research.

Figure 1: Summary of 3-digit octal games.

Code	$\mathcal{Q}$	p	Φ
0.34	$\mathcal{S}_{12}$	8	$a\ 1\ a\ b\ 1\ a\ 1\ ab\ a\ c\ a\ b\ 1\ ac\ 1\ ab\ a\ c\ a\ b\ 1\ ac\ \dots$
0.750	$\mathcal{R}_8$	2	$a\ b\ a\ b\ c\ b\ c\ b\ ab^2\ b\ ab^2\ b\ ab^2\ \dots$
0.512	$\mathcal{R}_8$	6	$a\ a\ a\ b\ b\ b\ a\ b^2\ c\ b\ b\ b\ ab^2\ b^2\ c\ b\ b\ b\ ab^2\ b^2\ ab^2\ b\ b\ b\ ab^2\ b^2\ ab^2\ \dots$
0.712	$\mathcal{R}_{14}$	6	$a\ b\ a\ 1\ c\ 1\ a\ ac\ a\ 1\ c\ 1\ a\ ac\ \dots$
0.716	$\mathcal{R}_{14}$	2	$a\ b\ a\ 1\ c\ 1\ c\ ac\ a\ ac\ c\ ac\ c\ ac\ c\ ac\ ac^2\ ac\ c\ c^2\ c\ c^2\ ac^2\ c^2\ ac^2\ c^2\ ac^2\ \dots$
4.56	$\mathcal{R}_8$	4	$a\ a\ b\ b\ a\ c\ b\ b\ c\ c\ b\ b\ ab^2\ ab^2\ b\ b\ ab^2\ ab^2\ \dots$
4.74	$\mathcal{R}_8$	2	$a\ b\ a\ b\ c\ b\ c\ b\ ab^2\ b\ ab^2\ b\ ab^2\ \dots$
0.3101	$\mathcal{R}_{14}$	2	$a\ b\ c^2\ c\ c^2\ ac^2\ c^2\ ac^2\ c^2\ ac^2\ \dots$
0.3131	$\mathcal{S}_{12}$	2	$a\ b\ a\ b\ b^2\ c\ b^2\ ab^2\ b^2\ ab^2\ b^2\ ab^2\ \dots$

Figure 2: Pretending functions for octal games with misère quotients isomorphic to $\mathcal{R}_8$, $\mathcal{S}_{12}$ or $\mathcal{R}_{14}$ (cf. Figure 10).

∞

Code	p	Φ	$\mathcal{Q}$	$\mathcal{P}$
0.1230	5	$a\ 1\ b\ b\ a\ d^2\ 1\ c\ d\ a\ d^2\ 1\ c\ d\ \dots$	$\langle a, b, c, d \mid a^2 = 1, b^4 = b^2,$ $b^2c = b^3, c^2 = 1, b^2d = d,$ $cd = bd, d^3 = ad^2 \rangle$	$\{a, b^2, ac,$ $bd, d^2\}$
0.1023	7	$a\ 1\ 1\ b\ b\ a\ a\ d^2\ 1\ 1\ c\ d\ a\ a\ d^2\ 1\ 1\ c\ d\ \dots$		
0.1032	7	$a\ 1\ a\ 1\ b\ b\ b\ a\ d^2\ d^2\ 1\ c\ ad^2\ d\ a\ d^2\ d^2\ 1\ c\ ad^2\ d\ \dots$		
0.1033	7	$a\ 1\ a\ b\ b\ b\ 1\ d^2\ d^2\ a\ d\ ad^2\ c\ 1\ d^2\ d^2\ a\ d\ ad^2\ c\ \dots$		
0.1331	5	$a\ a\ b\ b\ 1\ d^2\ a\ d\ c\ 1\ d^2\ a\ d\ c\ \dots$		
0.3103	5	$a\ b\ d^2\ d\ 1\ a\ c\ d^2\ d\ 1\ a\ c\ \dots$	$\langle a, b, c, d \mid a^2 = 1, b^4 = b^2,$ $b^2c = b^3, bc^3 = bc, c^4 = c^2,$ $bcd = b^2d, b^3d^2 = bd^2,$ $cd^2 = bd^2, b^2d^3 = d^3, d^5 = ad^4 \rangle$	$\{a, b^2, ac, ac^2,$ $ac^3, bd, ab^3d,$ $cd, ad^2, b^2d^2,$ $abd^3, d^4\}$
0.3112	5	$a\ b\ a\ 1\ b\ d^2\ c\ a\ 1\ d\ d^2\ c\ a\ 1\ d\ \dots$		
0.1232	6	$a\ 1\ b\ b\ ab\ a\ b^2\ 1\ c\ b\ d\ a\ b^2\ 1\ c\ b\ d\ \dots$		
0.1323	6	$a\ a\ 1\ b\ b\ ab\ a\ b^2\ 1\ c\ b\ d\ a\ b^2\ 1\ c\ b\ d\ \dots$		

Figure 3: Pretending functions and quotient presentations for some wild quaternary games.

Code	$\mathcal{Q}$	$\mathcal{P}$	Φ
0.530	$\langle a, b, c, d, e \mid a^2 = 1, b^3 = b, b^2c = c, c^2 = b^2, bd = ab, cd = ac, d^2 = b^2, be = ab, ce = ac, e^2 = b^2 \rangle$	$\{a, b^2, e\}$	$ \begin{array}{ccccccc} a & a & b & b & a & b^2 & b & b & c \\ b^2 & d & b & b & e & ab^2 & b & b & c \\ ab^2 & ade & b & b & ab^2 & ab^2 & b & b & c \\ ab^2 & ab^2 & b & b & ab^2 & ab^2 & b & b & c \\ ab^2 & ab^2 & \dots & & & & & & \end{array} $
0.710	$\langle a, b, c, d \mid a^2 = 1, b^4 = b^2, b^2c = c, c^4 = ac^3, c^3d = c^3, d^2 = 1 \rangle$	$\{a, b^2, bc, c^2, ac^3, ad, b^3d, cd, bc^2d\}$	$ \begin{array}{ccccccc} a & b & a & 1 & c & 1 & \\ a & d & a & 1 & c & 1 & \\ a & d & \dots & & & & \end{array} $
0.720	$\langle a, b, c, d, e, f \mid a^2 = 1, b^4 = b^2, b^2c = ab^2, c^2 = b^2, b^2d = b^3, cd = ab^3, d^2 = b^2, b^2e = ab^3, bce = bd, de = abd, e^2 = b^2, bf = bd, cf = ab^3, df = b^2, ef = ab^2, f^2 = b^2 \rangle$	$\{a, b^2, ad, be, af\}$	$ \begin{array}{ccccccc} a & 1 & b & ab & & & \\ a & 1 & b & ab & & & \\ c & b^2 & abc & ab^3 & & & \\ c & b^2 & d & e & & & \\ abd & b^2 & f & e & & & \\ abd & b^2 & f & \dots & & & \end{array} $
0.144	$\langle a, b, c, d, e \mid a^2 = 1, b^3 = b, b^2c = c, c^4 = c^2, bd = ab, cd = ac, d^2 = b^2, c^2e = c^3, de = ab^2e, e^2 = c^2 \rangle$	$\{a, b^2, c^2, ae, abce\}$	$ \begin{array}{cccccccc} a & 1 & 1 & a & b & b & b & c & c \\ a & ab^2 & ab^2 & d & b & b & b & b & c^3 & e \\ ab^2 & ab^2 & ab^2 & d & b & b & b & b & c^3 & e \\ ab^2 & \dots & & & & & & & & \end{array} $
0.241	$\cong \mathcal{Q}(\mathbf{0.71})$		$ \begin{array}{ccccccc} 1 & a & b & a & 1 & a & 1 & c & 1 & a \\ 1 & a & d & a & 1 & a & 1 & c & 1 & a \\ 1 & a & d & \dots & & & & & & \end{array} $
0.351	$\langle a, b, c, d \mid a^2 = 1, b^3 = b, b^2c = c, c^3 = ac^2, b^2d = d, c^2d = ac^2, d^2 = b^2 \rangle$	$\{a, b^2, bc, c^2, d, bcd\}$	$ \begin{array}{ccccccc} a & b & a & b & b^2 & c & b^2 & b \\ d & b & d & b & b^2 & c & b^2 & b \\ d & b & d & \dots & & & & \end{array} $

Figure 4: Quotient presentations and pretending functions for **0.53**, **0.71**, **0.72**, **0.144**, **0.241**, and **0.351**.

$$\mathcal{Q}(\mathbf{0.115}) \cong \langle a, b, c, d, e, f, g, h, i \mid a^2 = 1, b^4 = b^2, bc = ab^3, c^2 = b^2, \\ b^2d = d, cd = ad, d^3 = ad^2, b^2e = b^3, de = bd, be^2 = ace, \\ ce^2 = abe, e^4 = e^2, bf = b^3, df = d, ef = ace, cf^2 = cf, \\ f^3 = f^2, b^2g = b^3, cg = ab^3, dg = bd, eg = be, fg = b^3, \\ g^2 = bg, bh = bg, ch = ab^3, dh = bd, eh = bg, fh = b^3, \\ gh = bg, h^2 = b^2, bi = bg, ci = ab^3, di = bd, ei = be, \\ fi = b^3, gi = bg, hi = b^2, i^2 = b^2 \rangle$$

$$\mathcal{P} = \{a, b^2, bd, d^2, ae, ae^2, ae^3, af, af^2, ag, ah, ai\}$$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
0+	a	a	a	1	a	a	a	b	b	b	a	b	b	b
14+	a	a	a	1	c	c	c	b	b	b	d	b	b	e
28+	c	c	c	f	c	c	c	b	b	g	d	h	h	i
42+	ab^2	ab^2	abg	f	abg	abg	abe	b^3	b^3	h	d	h	h	h
56+	ab^2	ab^2	abe	f^2	abg	abg	abg	b^3	b^3	h	d	h	h	h
70+	ab^2	ab^2	abg	f^2	abg	abg	abg	b^3	b^3	b^3	d	b^3	b^3	b^3
84+	ab^2	ab^2	abg	f^2	abg	abg	abg	b^3	b^3	b^3	d	b^3	b^3	b^3
98+	ab^2	ab^2	ab^2	f^2	ab^2	ab^2	ab^2	b^3	b^3	b^3	d	b^3	b^3	b^3
112+	ab^2	ab^2	ab^2	f^2	ab^2	ab^2	ab^2	...						

Figure 5: Quotient presentation and pretending function for **0.115**.

$$\mathcal{Q}(\mathbf{0.152}) \cong \langle a, b, c, d, e \mid a^2 = 1, b^3 = b, bc^2 = b, c^3 = c, bd = b, c^2d = d, \\ d^2 = b^2, b^2e = e, c^2e = e, de = e, e^3 = ce^2 \rangle$$

$$\mathcal{P} = \{a, b^2, c^2, acd, abce, e^2\}$$

	1	2	3	4	5	6	7	8	9	10	11	12
0+	a	a	1	b	b	b	1	a	b^2	c	ab	b
12+	ab	ab^2	b^2	a	ab	b	b	ce^2	d	ab^2	b^2	e
24+	b	b	ab	ab^2	b^2	ab^2	ab	b	ab	ce^2	b^2	ab^2
36+	b^2	b	b	b	b^2	ab^2	b^2	ce^2	ab	b	ab	ce^2
48+	b^2	ab^2	b^2	b	b	b	b^2	ab^2	b^2	ce^2	ab	b
60+	ab	ab^2	b^2	ab^2	ab	b	b	ce^2	b^2	ab^2	b^2	ce^2
72+	b	b	ab	ab^2		...						

Figure 6: Quotient presentation and pretending function for **0.152**.

$$\mathcal{Q}(\mathbf{0.77}) \cong \langle a, b, c, d, e, f, g \mid a^2 = 1, b^3 = b, bc^2 = b, c^3 = c, bd = bc, \\ cd = b^2, d^3 = d, be = bc, ce = b^2, e^2 = de, \\ bf = ab, cf = ab^2c, d^2f = f, f^2 = b^2, b^2g = g, \\ c^2g = g, dg = cg, eg = cg, fg = ag, g^2 = b^2 \rangle$$

$$\mathcal{P} = \{a, b^2, ac, ac^2, d, ad^2, e, ade, adf\}$$

	1	2	3	4	5	6	7	8	9	10	11	12
0+	a	b	ab	a	c	ab	b	ab^2	d	b	bc	e
12+	ab^2	b	abc	ab^2	d^2e	ab	b	ade	b^2c	bc	abc	b^2c
24+	f	b	g	ab^2c	b^2c	abc	b	ab^2	g	bc	abc	b^2c
36+	ab^2	b	ab	ab^2	b^2c	abc	b	ab^2	g	b	abc	b^2c
48+	ab^2	b	g	ab^2	b^2c	abc	b	ab^2	b^2c	b	abc	b^2c
60+	ab^2	b	g	ab^2	b^2c	abc	b	ab^2	g	bc	abc	b^2c
72+	ab^2	b	g	ab^2	b^2c	abc	b	ab^2	g	b	abc	b^2c
84+	ab^2	b	g	ab^2	b^2c	abc	b	ab^2	g	b	$\dots$	

Figure 7: Quotient presentation and pretending function for **0.77**.

$$\mathcal{Q} \cong \langle a, b, c_n \mid a^2 = 1, b^{n+1}c_n = b^{2n+3}, \\ (c_m c_n = b^{m+2}c_n)_{m \leq n} \rangle$$

$$\mathcal{P} = \{a, (b^{2m})_{m \geq 1}, (b^m c_n)_{m \leq n \text{ and } m+n \text{ odd}}\}$$

	1	2	3	4
0+	1	a	b	ab
4+	1	a	b	ab
8+	c_0	ac_0	c_1	ab^3
12+	c_2	abc_1	c_3	abc_2
16+	c_4	abc_3	c_5	abc_4
20+	c_6	abc_5	$\dots$	

Figure 8: Quotient presentation and pretending function for **0.26**.

$$\mathcal{Q} \cong \langle a, b, c, d_n \mid a^2 = 1, bc = ab^3, c^2 = b^4, \\ b^{n+1}d_n = a^{n+1}b^{2n+5}, cd_n = ab^2d_n, \\ d_m d_n = a^{m+1}b^{m+4}d_n \rangle$$

$$\mathcal{P} = \{a, (b^{2m})_{m \geq 1}, (b^m d_n)_{m \text{ odd}, n \text{ even}, m < n}, \\ (ab^m d_n)_{m \text{ even}, n \text{ odd}, m < n}\}$$

	1	2
0+	a	b
2+	a	b
4+	c	b^3
6+	d_0	d_1
8+	d_2	d_3
10+	d_4	d_5
12+	d_6	$\dots$

Figure 9: Quotient presentation and pretending function for **4.7**.

Algebraic Periodic Games: 0.26 and 4.7

In Section A.5 we gave two examples of algebraic periodic games: **0.26** and **4.7** have infinite (and non-finitely generated) misère quotients. Furthermore, all of their *partial* quotients are finite. Full presentations are shown in Figures 8 and 9, respectively.

Since we do not have any computational methods for verifying algebraic periodicity, we must resort to manual proofs of Figures 8 and 9. In each case, the proof proceeds in two stages. We first show that the given presentation is *correct*, in the sense that G is a $\mathcal{P}$ -position iff $\Phi(G) \in \mathcal{P}$. Then we show that the given presentation is *reduced* (as a bipartite monoid).

Figure 8 Is Correct for 0.26

Let $\mathcal{A}$ be the set of **0.26** positions, regarded as a free commutative monoid on the heap alphabet $\mathcal{H} = \{H_1, H_2, H_3, \dots\}$. Define homomorphisms $t, w : \mathcal{A} \rightarrow \mathbb{N}$ as follows:

$$t(H_k) = \begin{cases} 0 & \text{if } k \in \{1, 2\}; \\ 1 & \text{if } k \in \{3, 4\}; \\ \lfloor \frac{k-5}{2} \rfloor & \text{if } k \geq 5. \end{cases} \quad w(H_k) = \begin{cases} 0 & \text{if } k \leq 8; \\ 1 & \text{if } k \in \{9, 10\}; \\ (k-7)/2 & \text{if } k \geq 11 \text{ is odd}; \\ (k-12)/2 & \text{if } k \geq 11 \text{ is even}. \end{cases}$$

Let $g(G)$ denote the ordinary Grundy value of G . It is easily checked that $g(H_k)$ is equal to the mod-4 parity of $k-1$.

Fact B.1 (Allemang [1]). *Let $G \in \mathcal{A}$ and write $G = X + H$, where $H = H_k$ is the largest single heap appearing in G . Then G is a $\mathcal{P}$ -position iff one of the following conditions holds:*

- (i) $t(G) = 0$ and $g(G) = 1$;
- (ii) $t(G) \neq 0$, $w(H) \leq t(X)$, and $g(G) = 0$; or
- (iii) $t(G) \neq 0$, $w(H) \geq t(X) + 2$, and $g(G) = 2$.

Fact B.1 corrects some slight errors in Allemang's definition of t . This would be a good reason to include a proof here; but the only proof we know is a tedious and unenlightening combinatorial slog, and anyway it is our hope that algorithmic verification methods will emerge in the near future. We therefore choose to take the easy way out, and leave its proof as an exercise.

Theorem B.2. *Figure 8 is correct for 0.26.*

Proof. Let $G \in \mathcal{A}$. We must show that G is a $\mathcal{P}$ -position iff $\Phi(G) \in \mathcal{P}$. We first make two preliminary observations, which are easily proved by inspecting the definitions of g , t , and Φ :

$$\text{If } k \leq k', \text{ then } \Phi(H_k)\Phi(H_{k'}) = a^{g(H_k)}b^{t(H_k)}\Phi(H_{k'}). \quad (\dagger)$$

$$\text{If } w(H) \leq m, \text{ then } b^m \Phi(H) = a^{g(H)} b^{m+t(H)}. \quad (\ddagger)$$

Now fix G , and write $G = X + H$ where $H = H_k$ is the largest heap appearing in G . By repeated application of $(\ddagger)$, we have $\Phi(G) = a^{g(X)} b^{t(X)} \Phi(H)$. There are five cases.

Case 1: $t(G) = 0$. Then $\Phi(G) = a^{g(G)}$ and the conclusion is evident.

case 2: $t(G) \neq 0$ and $w(H) \leq t(X)$. Then by $(\ddagger)$ we have

$$\Phi(G) = a^{g(X)} b^{t(X)} \Phi(H) = a^{g(X)} a^{g(H)} b^{t(X)+t(H)} = a^{g(G)} b^{t(G)}.$$

So $\Phi(G) \in \mathcal{P}$ iff $g(G)$ and $t(G)$ are both even. But $t(G)$ is even iff $g(G) \in \{0, 1\}$, so it follows that

$$\Phi(G) \in \mathcal{P} \text{ iff } g(G) = 0.$$

This agrees with the characterization in Fact B.1.

Case 3: $t(G) \neq 0$, $w(H) > t(X)$, and $k \in \{9, 10\}$. Then $w(H) = 1$, so $t(X) = 0$, and we have $\Phi(G) = a^{g(X)} \Phi(H)$. Since $\Phi(H_9) = c_0$ and $\Phi(H_{10}) = ac_0$, this necessarily implies $\Phi(G) \notin \mathcal{P}$, which agrees with Fact B.1.

Case 4: $t(G) \neq 0$, $w(H) > t(X)$, and $k \geq 11$ is odd. Then $\Phi(H) = c_{w(H)-1}$ and we have

$$\Phi(G) = a^{g(G)} b^{t(X)} c_{w(H)-1}.$$

Thus $\Phi(G) \in \mathcal{P}$ precisely when $g(G)$ and $t(X) + w(H)$ are both even. But $t(X) + w(H)$ is even iff $t(G)$ is odd, so this agrees with Fact B.1.

Case 5: $t(G) \neq 0$, $w(H) > t(X)$, and $k \geq 11$ is even. Then $\Phi(H) = abc_{w(H)}$ and so

$$\Phi(G) = a^{g(G)} b^{t(X)+1} c_{w(H)}.$$

Thus $\Phi(G) \in \mathcal{P}$ precisely when $g(G)$ and $t(X) + w(H)$ are both even, and the conclusion is just as in Case 4. $\square$

This shows that $\Phi : \mathcal{A} \rightarrow \mathcal{Q}$ is a homomorphism of bipartite monoids. The following theorem completes the picture.

Theorem B.3. *The bipartite monoid $(\mathcal{Q}, \mathcal{P})$ given in Figure 8 is reduced.*

Proof. For convenience, write $c_\infty = 1$. Then every $x \in \mathcal{Q}$ may be written uniquely as $a^i b^m c_n$, where $i \in \{0, 1\}$, $m \in \mathbb{N}$, $n \in \mathbb{N} \cup \{\infty\}$, and $m \leq n$. We will prove that, if $(i, m, n) \neq (i', m', n')$, then $x = a^i b^m c_n$ and $x' = a^{i'} b^{m'} c_{n'}$ are distinguishable.

First suppose $i \neq i'$, so without loss of generality $i = 0$ and $i' = 1$. Fix an integer $M > \max\{n, n'\}$ such that $m+n+M$ is even. Then $xb^M = b^{m+n+2+M} \in \mathcal{P}$, while $x'b^M = ab^{m'+n'+2+M} \notin \mathcal{P}$.

Now assume $i = i'$. It suffices to assume $i = 0$, since if x and x' are distinguished by z , then ax and ax' are distinguished by az . We may also assume, without loss of generality, that $n - m \leq n' - m'$. There are several cases.

Case 1: $m + n \not\equiv m' + n' \pmod{2}$. Then let $M > \max\{n, n'\}$ be such that $m + n + M$ is even. As before, $(b^m c_n) b^M = b^{m+n+2+M}$, while $(b^{m'} c_{n'}) b^M = b^{m'+n'+2+M}$. Since $m + n + 2 + M$ is even and $m' + n' + 2 + M$ is odd, we are done.

Case 2: $m + n \equiv m' + n' \pmod{2}$ and $n - m < n' - m'$. Without loss of generality, we may assume that $n - m < n' - m'$. Then $x b^{n-m+1} = b^{n+1} c_n = b^{2n+3} \notin \mathcal{P}$, while $x' b^{n'-m'+1} = b^{m'+n-m+1} c_{n'}$. But $m' + n - m + 1 < m' + n' - m' + 1 = n' + 1$, and

$$m' + n - m + 1 + n' \equiv (m' + n') + (n - m) + 1 \equiv (m' + n') + (m + n) + 1 \equiv 1 \pmod{2},$$

so this is in $\mathcal{P}$.

Case 3: $n - m = n' - m'$. Then we may assume that $n < n'$. Let $M > \max\{n, n'\}$ and put $N = m' + n' + 2 + M$. Then $x b^M c_N = b^{m'+n'+2+M} c_N \notin \mathcal{P}$. However,

$$x b^M c_N = b^{m+n+2+M} c_N = b^{m+n+4+M+N}.$$

Since $m + n \equiv m' + n' \pmod{2}$, this exponent is even. Hence $x b^M c_N \in \mathcal{P}$. $\square$

Figure 9 Is Correct for 4.7

Let $\mathcal{A}$ be the set of 4.7 positions, regarded as a free commutative monoid on the heap alphabet $\mathcal{H} = \{H_1, H_2, H_3, \dots\}$. Define homomorphisms $t, w : \mathcal{A} \rightarrow \mathbb{N}$ as follows:

$$t(H_k) = \begin{cases} 0 & \text{if } k = 1; \\ 1 & \text{if } k = 2; \\ k - 3 & \text{if } k \geq 3. \end{cases} \quad w(H_k) = \begin{cases} 0 & \text{if } k \leq 4; \\ 1 & \text{if } k = 5; \\ k - 6 & \text{if } k \geq 6. \end{cases}$$

Let $g(G)$ denote the ordinary Grundy value of G . It is easily checked that $g(H_k) = 2$ if k is even, 1 if k is odd.

Fact B.4 (Allemang [1]). *Let $G \in \mathcal{A}$ and write $G = X + H$, where $H = H_k$ is the largest single heap appearing in G . Then G is a $\mathcal{P}$ -position iff one of the following conditions holds:*

- (i) $t(G) = 0$ and $g(G) = 1$;
- (ii) $t(G) \neq 0$, $w(H) \leq t(X)$, and $g(G) = 0$; or
- (iii) $t(G) \neq 0$, $w(H) \geq t(X) + 2$, and $g(G) = 3$.

Theorem B.5. *Figure 9 is correct for 4.7.*

Proof. Fix $G \in \mathcal{A}$. We must show that G is a $\mathcal{P}$ -position iff $\Phi(G) \in \mathcal{P}$. As in the case of 0.26, the following observations are easily checked:

$$\text{If } k \leq k', \text{ then } \Phi(H_k) \Phi(H_{k'}) = a^{g(H_k)} b^{t(H_k)} \Phi(H_{k'}). \quad (\dagger)$$

If $w(H) \leq m$, then $b^m \Phi(H) = a^{g(H)} b^{m+t(H)}$. (†)

Now fix G , and write $G = X + H$ where $H = H_k$ is the largest heap appearing in G . By repeated application of (†), we have $\Phi(G) = a^{g(X)} b^{t(X)} \Phi(H)$. There are four cases.

Case 1: $t(G) = 0$. Then $\Phi(G) = a^{g(G)}$ and the conclusion is evident.

Case 2: $t(G) \neq 0$ and $w(H) \leq t(X)$. Then by (†) we have

$$\Phi(G) = a^{g(X)} b^{t(X)} \Phi(H) = a^{g(X)} a^{g(H)} b^{t(X)+t(H)} = a^{g(G)} b^{t(G)}.$$

So $\Phi(G) \in \mathcal{P}$ iff $g(G)$ and $t(G)$ are both even. But $t(G)$ is even iff $g(G) \in \{0, 1\}$, so this agrees with the characterization in Fact B.4.

Case 3: $t(G) \neq 0$, $w(H) > t(X)$, and $k \in \{5, 6\}$. Then $w(H) = 1$, so $t(X) = 0$, and we have $\Phi(G) = a^{g(X)} \Phi(H)$. Since $\Phi(H_5) = c$ and $\Phi(H_6) = b^3$, this necessarily implies $\Phi(G) \notin \mathcal{P}$, which agrees with Fact B.4.

Case 4: $t(G) \neq 0$, $w(H) > t(X)$, and $k \geq 7$. Then $\Phi(H) = d_{w(H)-1}$, so

$$\Phi(G) = a^{g(X)} b^{t(X)} d_{w(H)-1}.$$

Now if $w(H)$ is odd, then $\Phi(G) \in \mathcal{P}$ iff $g(X)$ is even and $t(X)$ is odd. But $w(H) \equiv g(H) \pmod{2}$, and $w(H) \not\equiv t(H) \pmod{2}$, so this means $g(G)$ and $t(G)$ are both *odd*. Since $t(G)$ is odd iff $g(G) \in \{2, 3\}$, this agrees with the characterization in Fact B.4.

Likewise, if $w(H)$ is even, then $\Phi(G) \in \mathcal{P}$ iff $g(X)$ is odd and $t(X)$ is even. Once again, this means that $g(G)$ and $t(G)$ are both odd. This exhausts all cases and completes the proof. $\square$

Theorem B.6. *The bipartite monoid $(\mathcal{Q}, \mathcal{P})$ given in Figure 9 is reduced.*

The proof of Theorem B.6 is much like Theorem B.3.

Games Born by Day 4

The quotients $\mathcal{Q}(G)$, for a single misère game G , are fundamental (but in some cases still quite intricate). Figure 10 summarizes all quotients obtained this way, to birthday four. Each quotient is listed together with its order, monoid presentation, and $\mathcal{P}$ -portion.

The canonical forms of the twenty-two misère impartial games G of birthday at most four were introduced by Conway [2], and are duplicated here in Figure 11. They are shown together with their misère quotients.

There are exactly 4171780 games born by day five [2]. They yield a bewildering variety of misère quotients, including infinite quotients; finite quotients with more than 1500 elements (and probably larger ones as well); and counterexamples to several reasonable-sounding statements about the general structure of misère quotients. It is conceivable that a complete survey of games born by day five might eventually be conducted.

	$ \mathcal{Q} $	$\mathcal{Q}$	$\mathcal{P}$
$\mathcal{T}_0$	1	$\langle \mid \rangle$	$\emptyset$
$\mathcal{T}_1$	2	$\langle a \mid a^2 = 1 \rangle$	$\{a\}$
$\mathcal{T}_2$	6	$\langle a, b \mid a^2 = 1, b^3 = b \rangle$	$\{a, b^2\}$
$\mathcal{R}_8$	8	$\langle a, b, c \mid a^2 = 1, b^3 = b, bc = ab, c^2 = b^2 \rangle$	$\{a, b^2\}$
$\mathcal{T}_3$	10	$\langle a, b, c \mid a^2 = 1, b^3 = b, c^3 = c, c^2 = b^2 \rangle$	$\{a, b^2\}$
$\mathcal{S}_{12}$	12	$\langle a, b, c \mid a^2 = 1, b^4 = b^2, b^2c = b^3, c^2 = 1 \rangle$	$\{a, b^2, ac\}$
$\mathcal{S}'_{12}$	12	$\langle a, b, c \mid a^2 = 1, b^3 = b, c^2 = 1 \rangle$	$\{a, b^2, ac\}$
$\mathcal{R}_{14}$	14	$\langle a, b, c \mid a^2 = 1, b^3 = b, b^2c = c, c^3 = ac^2 \rangle$	$\{a, b^2, bc, c^2\}$

Figure 10: The eight misère quotients born by day 4.

0	*	*2 *3/ *2/2	*3 *32 *2/210 *2/32	*2/ *2// *2/3	*2/320	*4 *2/3210	*2/1 *2/21 *2/31	*2/321	*2/0 *2/20 *2/30
$\mathcal{T}_0$	$\mathcal{T}_1$	$\mathcal{T}_2$			$\mathcal{R}_8$	$\mathcal{T}_3$	$\mathcal{S}_{12}$	$\mathcal{S}'_{12}$	$\mathcal{R}_{14}$

Figure 11: The 22 misère games born by day 4, grouped by quotient isomorphism type.

The Games $0.(\mathbf{3310})^n$

Suppose misère Nim is played with two restrictions:

- $4k$ beans may not be removed from any heap;
- $4k + 3$ beans may be removed only if it is the whole heap.

In standard notation, this game would be

$$0.(\mathbf{3310})^\infty = \mathbf{0.3310331033103310} \dots$$

Does its misère quotient have order 226? The following approximations seem to suggest that it does:

Γ	$ \mathcal{Q} $	pd
0.3310	6	3
0.33103310	202	7
0.331033103310	226	11
0.3310331033103310	226	15
0.33103310331033103310	226	19

Question. Does $\mathcal{Q}(0.(\mathbf{3310})^\infty) = \mathcal{Q}(0.(\mathbf{3310})^n)$ for all $n \geq 3$? Can we say anything interesting in general about games with infinite octal codes?

C Algorithms

In this appendix we describe *MisereSolver*'s algorithm for calculating misère quotients. Throughout this section, let $\mathcal{A}$ be an arbitrary finitely-generated closed set of games, treated as a free commutative monoid on generators $\mathcal{H} = \{H_1, H_2, \dots, H_n\}$. Denote by $<_{\text{lex}}$ the lexicographic ordering on $\mathcal{A}$:

$$\left(\sum k_i H_i\right) <_{\text{lex}} \left(\sum \ell_i H_i\right) \text{ iff } k_i < \ell_i \text{ for the largest } i \text{ such that } k_i \neq \ell_i.$$

Notice that if $X \in \mathcal{A}$ and X' is an option of X , then $X' <_{\text{lex}} X$.

The basic idea is to produce a sequence of increasingly accurate approximations to the misère quotient $\mathcal{Q}(\mathcal{A})$. If $\mathcal{Q}(\mathcal{A})$ is indeed finite, then the sequence is guaranteed to converge to it.

At each stage of the iteration, we are given a promising r.b.m. $(\mathcal{Q}, \mathcal{P})$ (the “candidate quotient”), together with a homomorphism $\Phi : \mathcal{A} \rightarrow \mathcal{Q}$ (represented as a mapping $\mathcal{H} \rightarrow \mathcal{Q}$). If $(\mathcal{Q}, \mathcal{P})$ is the correct quotient of $\mathcal{A}$, with pretending function Φ , then we are done. If not, then there is some $X \in \mathcal{A}$, $X \neq 0$, such that either:

- (i) $\Phi(X) \in \mathcal{P}$, but also $\Phi(X') \in \mathcal{P}$ for some option X' ; or
- (ii) $\Phi(X) \notin \mathcal{P}$, but there is no option X' with $\Phi(X') \in \mathcal{P}$.

We say that such an X is a *failure* of the candidate $(\mathcal{Q}, \mathcal{P}, \Phi)$. Now suppose that X is the *lexicographically least* failure of $(\mathcal{Q}, \mathcal{P}, \Phi)$. Then, by a straightforward induction, Φ correctly predicts the outcomes of all proper followers of X . Therefore we know the outcome of X : it is necessarily the opposite of that predicted by Φ . Using this information, we produce a new candidate $(\mathcal{Q}', \mathcal{P}', \Phi')$, for which X is *not* a failure. The construction will guarantee that no lexicographically smaller failures are introduced. The least failures of successive candidate quotients are therefore strictly lexicographically increasing.

There are two main components of this algorithm.

- *Verification*: Given a candidate $(\mathcal{Q}, \mathcal{P}, \Phi)$, the verification engine determines whether it is the correct quotient of $\mathcal{A}$, and efficiently identifies the least failure when it is not.
- *Recalibration*: Given a candidate $(\mathcal{Q}, \mathcal{P}, \Phi)$, together with the least failure X , the recalibration engine constructs the next candidate $(\mathcal{Q}', \mathcal{P}', \Phi')$.

When *MisereSolver* begins, it starts with an initial round of recalibration. For the initial candidate, we use the misère quotient of $\langle H_1, \dots, H_{n-1} \rangle$ (computed recursively), taking $X = H_n$ to be the least failure.

We are reasonably certain that *MisereSolver* uses the “book” algorithm for verification: it is both elegant and fast. (In particular, we have improved substantially on the algorithms described in [3].) However, our recalibration algorithm feels very crude: *MisereSolver* sometimes chooses very poor candidates; often they are considerably larger than the true quotient. An improved recalibrator might dramatically extend *MisereSolver*'s scope.

Verification

Given a r.b.m. $(\mathcal{Q}, \mathcal{P})$ and a monoid homomorphism $\Phi : \mathcal{A} \rightarrow \mathcal{Q}$, we wish to determine:

- (a) The $<_{\text{lex}}$ -least $X \in \mathcal{A}$ such that $\Phi(X) \in \mathcal{P}$ but $\Phi(X') \notin \mathcal{P}$ for some option X' ;
- (b) The $<_{\text{lex}}$ -least $X \in \mathcal{A}$ such that $\Phi(X) \notin \mathcal{P}$ but $\Phi(X') \notin \mathcal{P}$ for all options X' .

As observed in [3], (b) is far more difficult computationally. This is due to the presence of the universal quantifier: in searching for *some* option in (a), we can essentially treat each generator as an independent entity; but to iterate over *all* options in (b), we must consider the totality of generators involved in X .

For (a), we use the essential strategy outlined in [3]. The basis is the following theorem:

Fact C.1 ($\mathcal{P}$ -Verification Theorem [3, Section 9.1.3]). *Let $(\mathcal{Q}, \mathcal{P})$ be a r.b.m. and fix a homomorphism $\Phi : \mathcal{A} \rightarrow \mathcal{Q}$. The following are equivalent.*

- (i) *There is a game $X \in \mathcal{A}$ and an option X' such that $\Phi(X), \Phi(X') \in \mathcal{P}$.*
- (ii) *There is a generator $H \in \mathcal{H}$, an option H' , and an element $x \in \mathcal{Q}$ such that $x\Phi(H), x\Phi(H') \in \mathcal{P}$.*

Therefore, to resolve (a), we can iterate over all pairs $(H, x) \in \mathcal{H} \times \mathcal{Q}$, testing condition (ii) in Fact C.1. For each such pair satisfying (ii), let X be the $<_{\text{lex}}$ -least game with $\Phi(X) = x$; then $H + X$ is a candidate for the $<_{\text{lex}}$ -least failure. This yields at most $|\mathcal{H}| \cdot |\mathcal{Q}|$ candidates, over which we simply minimize.

Our technique for (b) uses the following central idea.

Definition C.2. *Let $X, Y \in \mathcal{A}$, and suppose that $X <_{\text{lex}} Y$, $\Phi(X) = \Phi(Y)$, and $\Phi''X \subset \Phi''Y$. Then we say that Y is subsumed by X .*

Now if Y is subsumed by X , then Y cannot be the $<_{\text{lex}}$ -least failure of $(\mathcal{Q}, \mathcal{P}, \Phi)$, because if Y satisfies (b), then so does X . Moreover, $Y + Z$ cannot be the $<_{\text{lex}}$ -least failure, for *any* $Z \in \mathcal{A}$: once again, if $Y + Z$ satisfies (b), then so does $X + Z$; and necessarily $X + Z <_{\text{lex}} Y + Z$. Therefore, we can completely disregard any element of $\mathcal{A}$ containing Y as a subword.

We may therefore traverse the set $\mathcal{A}$ in lexicographic order, pruning whenever we reach an $X \in \mathcal{A}$ that we know to be subsumed. Since $\mathcal{Q}$ is finite and there are at most $|\mathcal{Q}| \cdot 2^{|\mathcal{Q}|}$ possibilities for $(\Phi(X), \Phi''X)$, this traversal is guaranteed to terminate (even though $\mathcal{A}$ is infinite). The full procedure is summarized as Algorithm 1.

```

1:  $T \leftarrow \emptyset$ 
2: return LEXLEASTNFAILURE( $n, 0$ )

3: procedure LEXLEASTNFAILURE( $d, X$ )
4:   if  $d = 0$  then                                ▷ Done recursing; check this value of  $X$ 
5:     if  $X = 0$  then
6:       return NULL                                ▷ 0 must be special-cased in misère play
7:     end if
8:      $x \leftarrow \Phi(X); \mathcal{E} \leftarrow \Phi''X$ 
9:     if  $x \in \mathcal{P}$  then
10:      return NULL                                ▷ We are seeking  $\mathcal{N}$ -failures only
11:    else
12:      if  $\mathcal{P} \cap \mathcal{E} = \emptyset$  then
13:        return  $X$                                 ▷ We've found the lex-least failure
14:      else
15:         $T \leftarrow T \cup \{(x, \mathcal{E})\}$           ▷ Add a transition record for  $(x, \mathcal{E})$ 
16:        return NULL                                ▷ Keep searching
17:      end if
18:    end if
19:  else                                              ▷ Continue recursing from depth  $d$ 
20:    loop
21:       $Y \leftarrow \text{LEXLEASTNFAILURE}(d - 1, X)$ 
22:      if  $Y \neq \text{NULL}$  then
23:        return  $Y$                                 ▷ Found the answer; return it
24:      end if
25:       $X \leftarrow X + H_d$                         ▷ Add a heap  $H_d$  to  $X$ 
26:       $x \leftarrow \Phi(X); \mathcal{E} \leftarrow \Phi''X$ 
27:      if there exists  $\mathcal{D} \subset \mathcal{E}$  with  $(x, \mathcal{D}) \in T$  then
28:        return NULL                                ▷  $X$  is subsumed; return immediately
29:      end if
30:    end loop
31:  end if
32: end procedure

```

Algorithm 1: $\mathcal{N}$ -Verification Algorithm.

Recalibration

Given an r.b.m. $(\mathcal{Q}, \mathcal{P})$, a map $\Phi : \mathcal{H} \rightarrow \mathcal{Q}$, and a lexicographically least failure $X \in \mathcal{A}$ (as certified by Verification), we must construct a new triple $(\mathcal{Q}', \mathcal{P}', \Phi')$ whose least failure Y is guaranteed to satisfy $Y >_{\text{lex}} X$. The construction must further guarantee that if $\mathcal{Q}(\mathcal{A})$ is finite, and if $(\mathcal{Q}_1, \mathcal{P}_1, \Phi_1), (\mathcal{Q}_2, \mathcal{P}_2, \Phi_2), \dots$ is a sequence of successive recalibrations, then $\mathcal{Q}(\mathcal{A}) = (\mathcal{Q}_n, \mathcal{P}_n)$ for sufficiently large n .

What follows is an informal description of *MisereSolver*'s recalibration procedure. The essential idea is to “free” all generators involved in the failure X , mapping them to suitable monoids of the form $\langle t \mid t^{n+k} = t^n \rangle$. This gives an expanded monoid, in which the true outcome of X may be “marked” without affecting the indicated outcome of any $Y <_{\text{lex}} X$. The expanded monoid is then immediately collapsed back down to a r.b.m. This ensures that we can recalibrate as many times as necessary while keeping the candidate quotients reasonably small.

Given a candidate $(\mathcal{Q}, \mathcal{P}, \Phi)$ with $<_{\text{lex}}$ -least failure X :

1. Let $\mathcal{J} = \{J_1, \dots, J_m\} \subset \mathcal{H}$ be the set of generators appearing in X . Let $\mathcal{Q}^-$ be the submonoid of $\mathcal{Q}$ generated by $\{\Phi(H) : H \in \mathcal{H} \setminus \mathcal{J}\}$. For each i ($1 \leq i \leq m$), let $\mathcal{R}_i = \langle t \mid t^{n+k} = t^n \rangle$, for some carefully chosen k, n .¹ Now put

$$\mathcal{Q}^* = \mathcal{Q}^- \times \mathcal{R}_1 \times \mathcal{R}_2 \times \dots \times \mathcal{R}_m.$$

2. Define $\Phi^* : \mathcal{H} \rightarrow \mathcal{Q}^*$ as follows. For each i , put $\Phi^*(J_i) =$ the generator of $\mathcal{R}_i$. For $H \notin \mathcal{J}$, put $\Phi^*(H) = (\Phi(H), 1, 1, \dots, 1)$.
3. Define $\mathcal{P}^* \subset \mathcal{Q}^*$ as follows. First put $1 \notin \mathcal{P}^*$. Then for each $x \in \mathcal{Q}^*$, let X be the $<_{\text{lex}}$ -least element of $\mathcal{A}$ with $\Phi^*(X) = x$. Since $X' <_{\text{lex}} X$ for each option X' of X , we may assume that we have already defined whether each $\Phi^*(X') \in \mathcal{P}^*$. So define

$$x \in \mathcal{P}^* \iff \Phi^*(X') \notin \mathcal{P}^* \text{ for each option } X'.$$

Essentially, we are guessing whether each such X is a $\mathcal{P}$ -position, under the assumption that Φ^* is correct on all its options.

4. This gives a bipartite monoid $(\mathcal{Q}^*, \mathcal{P}^*)$. Compute its reduction $(\mathcal{Q}', \mathcal{P}')$ and let Φ' be the factor map.

There is a great deal of inefficiency in this process: much more of the monoid is “freed” than is needed to enforce the correct outcome of X . Nonetheless, it works quite well for our purposes, and it satisfies the following two crucial theorems, which we state without proof.

¹*MisereSolver*'s method of choosing such k, n is currently rather crude and is not worth an extended discussion. The important thing is that k and n must be large enough to arrive eventually at the correct presentation, but small enough to keep the computations tractable. There is much room for improvement here.

Theorem C.3 (Recalibration Theorem). *Let $(\mathcal{Q}, \mathcal{P})$ be a r.b.m. and $\Phi : \mathcal{A} \rightarrow \mathcal{Q}$ a homomorphism with $<_{\text{lex}}$ -least failure X . Let $(\mathcal{Q}', \mathcal{P}', \Phi')$ be the recalibration of $(\mathcal{Q}, \mathcal{P}, \Phi)$ by X and let Y be the lex-least failure of Φ' . Then $X <_{\text{lex}} Y$.*

Theorem C.4 (Termination Theorem). *Let $\mathcal{A}$ be a closed set of games with finite misère quotient $(\mathcal{Q}, \mathcal{P})$ and pretending function $\Phi : \mathcal{A} \rightarrow \mathcal{Q}$. Let*

$$(\mathcal{Q}_1, \mathcal{P}_1, \Phi_1), (\mathcal{Q}_2, \mathcal{P}_2, \Phi_2), \dots$$

be a sequence of candidate quotients for $\mathcal{A}$, with each $(\mathcal{Q}_{n+1}, \mathcal{P}_{n+1}, \Phi_{n+1})$ computed by recalibrating on the $<_{\text{lex}}$ -least failure of $(\mathcal{Q}_n, \mathcal{P}_n, \Phi_n)$. Then for all sufficiently large n , we have $(\mathcal{Q}_n, \mathcal{P}_n, \Phi_n) = (\mathcal{Q}, \mathcal{P}, \Phi)$.

In the raw form described here, Recalibration can be extremely slow. If the least failure X involves many generators, the expanded monoid $\mathcal{Q}^*$ is often quite large. Fortunately, a tremendous shortcut is available: in step 1, rather than expand along the entire set $\mathcal{J}$, we can first try to expand along some proper subset $\mathcal{J}' \subsetneq \mathcal{J}$. This almost always succeeds—in many cases, when $\mathcal{J}'$ is a singleton.

Optimizations for Heap Games

MisereSolver's chief application is to calculate misère quotients of heap games. When Γ is an octal game, it is likely that $\mathcal{Q}_{n+1}(\Gamma) = \mathcal{Q}_n(\Gamma)$ for many values of n . In such cases *MisereSolver* uses significant optimizations to compute the next pretension $\Phi(H_{n+1})$.

First, whenever a new partial quotient $\mathcal{Q}_n(\Gamma)$ is computed, *MisereSolver* also computes the meximal sets $\mathcal{M}_x$ for each $x \in \mathcal{Q}_n$. It also computes the antichain of lower bounds of the transition algebra T_n . (See Section 5 for discussion.) The algorithm used to compute this antichain is virtually identical to Algorithm 1.

Then, before computing $\mathcal{Q}_{n+1}(\Gamma)$, *MisereSolver* first calculates the set $\mathcal{E} = \Phi''H_{n+1}$. If there exists an $x \in \mathcal{Q}_n$ and a lower bound $(x, \mathcal{D}) \in T_n$ such that $\mathcal{D} \subset \mathcal{E} \subset \mathcal{M}_x$, then by the Mex Interpolation Principle it follows that $\Phi(H_{n+1}) = x$. This check is extremely fast. Furthermore, by the strong form of the Mex Interpolation Principle, it is not necessary to update the information about T_n . Thus *MisereSolver* can run quickly through a large number of interpolated heaps.

If no such lower bound $(x, \mathcal{D})$ exists, it might still be the case that $\mathcal{Q}_{n+1}(\Gamma) = \mathcal{Q}_n(\Gamma)$. To test this, *MisereSolver* tries every x with $\mathcal{E} \subset \mathcal{M}_x$. For each such x , the software assigns $\Phi(H_{n+1}) = x$ and runs Algorithm 1. If the algorithm returns NULL for some value of x , then we are done. (However, we must recompute the antichain of lower bounds of T_{n+1} .) If it fails on every x , then *MisereSolver* proceeds with the full recalibration/verification procedure described above.

References

- [1] D. T. Allemang. Machine computation with finite games. Master's thesis, Trinity College, Cambridge, 1984.
<http://www.miseregames.org/allemang/>.
- [2] J. H. Conway. *On Numbers and Games*. A. K. Peters, Ltd., Natick, MA, second edition, 2001.
- [3] T. E. Plambeck. Taming the wild in impartial combinatorial games. *INTEGERS: The Electr. J. Combin. Number Thy.*, 5(#G05), 2005.
- [4] T. E. Plambeck and A. N. Siegel. Misère Games on the Web.
<http://www.miseregames.org/>.
- [5] T. E. Plambeck and A. N. Siegel. Misère quotients for impartial games. Forthcoming. <http://arxiv.org/abs/math.CO/0609825>.